




**INBIOBANK** [www.inbiobank.org](http://www.inbiobank.org)  
Paseo Mikeletegi, 61 bajo. San Sebastian 20009 (Spain)  
Phone: +34 943 309064 Fax: +34 943 308222  
info@inbiobank.org

## HUMAN BONE MARROW MESENCHYMAL STEM CELLS (hMSC)

### QUALITY CONTROL






All cells have been processed in INBIOBANK following manufacturing procedures based on ISO9001:2000 norm on white room laboratory. Each cell donor is tested and found negative for: HIV-1, HIV-2, Hepatitis B and C (Abbot Prism chemiluminiscence assay) and mycoplasma (PCR Kit).

Cellular morphology and viability have been tested after criopreservation.

### BIOLOGICAL CONTROL





Human Mesenchymal Stem Cells display a typical CD29+, CD73+ (SH3 and SH4), CD105+ (SH2), CD166+, CD45- and CD31- phenotype. In presence of specific differentiation factors, these cells have been shown to differentiate to osteocytes, chondrocytes and fatty cells. (Ref. 1,2).



**BMONT#45MSC1**

**AGE: 29 YEARS OLD**  
**SEX: MALE**

Filled area: "ISOTYPIC CONTROL"  
Empty area: "SPECIFIC CD"





**INBIOBANK** [www.inbiobank.org](http://www.inbiobank.org)  
Paseo Mikeletegi, 61 bajo. San Sebastian 20009 (Spain)  
Phone: +34 943 309064 Fax: +34 943 308222  
[info@inbiobank.org](mailto:info@inbiobank.org)

---

### **Instructions for use:**

#### **Thawing of cells/initiation of culture**

Frozen ampoules: These are 1 ml plastic cryotubes containing **1 million** cells and FBS (v/v) dimethylsulphoxide (DMSO). The cells are supplied in dry ice. When they are received, store in liquid nitrogen or use them immediately.

- Remove vial from Liquid Nitrogen or dry ice and immediately transfer to 37°C water bath.
- While holding the tip of the vial, gently agitate the vial, being careful not to allow water to penetrate the cap or seal.
- Add 10ml warm complete media to a 15ml tube. When completely thawed, slowly transfer contents of vial to 15ml test tube and spin at 300g for 5min
- Remove media and resuspend pellet in a volume of complete media appropriate for flask (T75 flask).

#### **Subculturing**

- INBIOBANK recommends to plate a density of 1000-2000 cells/cm<sup>2</sup> for human mesenchymal stem cells
- hMSC cultures should be fed 3-4 days after plating. To feed the cultures, gently remove the medium from the well and replace with an equal volume of temperature equilibrated fresh medium.
- The hMSC should be subcultured when they are confluent (approximately 90%).

#### **Culture media.**

- Use Low-Glucose DMEM (Dulbecco's modified eagle's medium) supplemented with 10% Foetal Bovine Serum to feed human mesenchymal stem cells. Users should test a specific batch of serum for optimum growth of these cells.

#### **SAFETY STAMENTS.**

THIS MATERIAL IS FOR RESEARCH USE ONLY. Do not use in human assays, transplant or other *in vitro* diagnosis.

#### **REFERENCES**

- 1.-Pittenger, M. F., A. M. Mackay, et al. (1999). "Multilineage potential of adult human mesenchymal stem cells." *Science* **284**(5411): 143-7.
2. Javazon, E. H., K. J. Beggs, et al. (2004). "Mesenchymal stem cells: paradoxes of passaging." *Exp Hematol* **32**(5): 414-25.

