




## PRIMARY HUMAN KERATINOCYTES

Processed directly from adult human skin (from healthy donors) and expanded *in vitro* for a maximum of three passages in the presence of a lethally irradiated 3T3 feeder cell layer, by following the classical procedure of Rheinwald and Green [1, 2]. Cells are supplied in 2 ml cryotubes containing **1 million** cells in 1 ml FBS (v/v) dimethylsulphoxide (DMSO).




### Quality Controls

All cells have been processed in INBIOBANK by manufacturing procedures based on ISO9001:2000 standards. These cells have tested negative for the following pathogens: HIV-1/2, Hepatitis B/C (Abbot Prism Chemiluminescence Assay) and Mycoplasma (PCR Kit).

Cellular morphology and viability have also been tested after recovery from criopreservation.


### Cell characterisation

Human keratinocytes are characterised morphologically through serial passage, and each batch is tested for expression of the following keratinocyte markers: cytokeratin 14 (K14), integrin alpha-6 (CD49f), integrin beta-1 (CD29).



S0701K  
Age 39  
Sex: Female

Red area: Isotypic control  
Green line: specific CD





**INBIOBANK** [www.inbiobank.org](http://www.inbiobank.org)  
Paseo Mikeletegi, 61 bajo. San Sebastian 20009 (Spain)  
Phone: +34 943 309064 Fax: +34 943 308222  
info@inbiobank.org

## Instructions for use

### Storage:

- The cells may be supplied criopreserved in dry ice. When received, liquid nitrogen storage is recommended until use.

### Cell thawing:

- Transfer to 37°C water bath.
- Gently agitate the vial while holding the tip with a forceps, being careful not to allow water to penetrate the cap or seal.
- Quickly sterilize vial with 70% ethanol and transfer to a laminar flow hood. Add 10ml warm complete media to a 15ml tube. When completely thawed, slowly transfer contents of vial to 15ml test tube and spin at 300g for 5min.
- Remove and discard the medium and resuspend in fresh culture medium (w/o Epidermal growth factor, EGF).

### Plating and passaging of keratinocyte cultures.

- Cells must be seeded simultaneously with lethally irradiated mouse 3T3 cells (Swiss Albino, ECACC #85022108). These cells are irradiated at 50 Gy and stored at 4°C until use for a maximum of 4 days.
- INBIOBANK recommends a plating density of about 25,000 keratinocytes/cm<sup>2</sup> and 90,000 low passage 3T3 cells/cm<sup>2</sup>, in medium w/o EGF.
- Do not disturb the plates for the first 48 hours.
- Change the culture medium to medium with EGF, and refresh media every 2-3 days. Feeder layer also needs replenishment every week.
- Expand the cells when 80-90% confluent.

## Warning

FOR RESEARCH USE ONLY. Not for use in clinical or veterinary assays, transplants or *in vitro* diagnostic procedures.

### Human keratinocyte growth medium

Dulbecco's modified Eagle's medium (DMEM, 4,500 mg glucose/l)  
Ham's F-12 Nutrient mixture

### Working conc.

60%  
30%





**INBIOBANK** [www.inbiobank.org](http://www.inbiobank.org)  
Paseo Mikeletegi, 61 bajo. San Sebastian 20009 (Spain)  
Phone: +34 943 309064 Fax: +34 943 308222  
[info@inbiobank.org](mailto:info@inbiobank.org)

---

Foetal bovine serum <sup>†</sup>	10%
L-Glutamine	584 mg/l
Penicillin/Streptomycin	50 ug/ml
Adenin	0,18 mM
Cholera toxin	0,1 nM
Thyroid hormone T3	2 nM
Hydrocortisone	0,4 ug/ml
Bovine insulin	5 ug/ml
Epidermal growth factor (EGF) <sup>‡</sup>	10 ng/ml

### References

1. Rheinwald, J.G. and H. Green, *Serial cultivation of strains of human epidermal keratinocytes: the formation of keratinizing colonies from single cells*. Cell, 1975. **6**(3): p. 331-43.
2. Rheinwald, J.G. and H. Green, *Epidermal growth factor and the multiplication of cultured human epidermal keratinocytes*. Nature, 1977. **265**(5593): p. 421-4.

---

<sup>†</sup> End user must test specific serum batches for optimal cell growth.

<sup>‡</sup> Media must be prepared with and w/o EGF.

